

Package leaflet: Information for the patient

Eslo Tan[®] Tablets

S(-) Amlodipine 2.5 mg + Losartan Potassium 50 mg tablets

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you:

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Eslo Tan[®] is and what it is used for
2. What you need to know before you take Eslo Tan[®]
3. How to take Eslo Tan[®]
4. Possible side effects
5. How to store Eslo Tan[®]
6. Contents of the pack and other information

1. What Eslo Tan[®] is and what it is used for

Eslo Tan[®] tablets contain two substances called S(-)Amlodipine and Losartan. Both of these substances help to control high blood pressure.

– S(-)Amlodipine belongs to a group of substances called “calcium channel blockers”. S(-)Amlodipine stops calcium from moving into the blood vessel wall which stops the blood vessels from tightening.

– Losartan belongs to a group of substances called “angiotensin-II receptor antagonists”. Angiotensin II is produced by the body and makes the blood vessels tighten, thus increasing the blood pressure. Losartan works by blocking the effect of angiotensin II.

This means that both substances help stop the blood vessel tightening. As a result, the blood vessels relax and blood pressure is lowered.

Eslo Tan[®] is used to treat high blood pressure in adults whose blood pressure is not controlled enough with either S(-)Amlodipine or Losartan on its own.

2. What you need to know before you take Eslo Tan[®]

Do not take Eslo Tan[®]:

- if you are allergic to S(-)-Amlodipine, amlodipine, Losartan, any drug of a related class, or any of the other ingredients of this medicine (listed in section 6).
- If you think you may be allergic, talk to your doctor before taking Eslo Tan[®].
- if you have severe liver problems or bile problems such as biliary cirrhosis or cholestasis.
- if you are more than 3 months pregnant. (It is also better to avoid Eslo Tan[®] in early pregnancy, see Pregnancy section).
- if you have severe low blood pressure (hypotension).
- if you have narrowing of the aortic valve (aortic stenosis) or cardiogenic shock (a condition where your heart is unable to supply enough blood to the body).
- if you suffer from heart failure after a heart attack.
- if you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren.

If any of the above applies to you, do not take Eslo Tan[®] and talk to your doctor.

Warnings and precautions

Talk to your doctor before taking Eslo Tan[®] Tablets:

- if you have been sick (vomiting or diarrhoea).
- if you have liver or kidney problems.
- if you have had a kidney transplant or if you had been told that you have a narrowing of your kidney arteries.
- if you have a condition affecting the renal glands called “primary hyperaldosteronism”.
- if you have had heart failure or have experienced a heart attack. Follow your doctor’s instructions for the starting dose carefully. Your doctor may also check your kidney function.
- if your doctor has told you that you have a narrowing of the valves in your heart (called “aortic or mitral stenosis”) or that the thickness of your heart muscle is abnormally increased (called “obstructive hypertrophic cardiomyopathy”).
- if you have experienced swelling, particularly of the face and throat while taking other medicines (including angiotensin-converting enzyme inhibitors). If you get these symptoms, stop taking Eslo Tan[®] and contact your doctor straight away. You should never take Eslo Tan[®] again.
- if you are taking any of the following medicines used to treat high blood pressure:
 - an ACE inhibitor (for example enalapril, lisinopril, ramipril), in particular if you have diabetes-related kidney problems.
 - aliskiren.

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals.

See also information under the heading “Do not take Eslo Tan[®]”.

If any of these apply to you, tell your doctor before taking Eslo Tan[®].

Children and adolescents

The use of Eslo Tan[®] in children and adolescents is not recommended (aged below 18 years old).

Other medicines and Eslo Tan[®] Tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. Your doctor may need to change your dose and/or to take other precautions. In some cases, you may have to stop taking one of the medicines. This applies especially to the medicines listed below:

Eslo Tan[®] Tablets may affect or be affected by other medicines, such as:

- ACE inhibitors or aliskiren (see also information under the headings “Do not take Eslo Tan[®]” and “Warnings and precautions”);
- diuretics (which increases the amount of urine you produce);
- lithium (a medicine used to treat some types of depression);
- potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium and other substances that may increase potassium levels;
- certain types of painkillers called non-steroidal anti-inflammatory medicines (NSAIDs) or selective cyclooxygenase-2 inhibitors (COX-2 inhibitors). Your doctor may also check your kidney function;
- anticonvulsant agents (e.g. carbamazepine, phenobarbital, phenytoin, fosphenytoin, primidone).
- St. John’s wort;
- nitroglycerin and other nitrates, or other substances called “vasodilators”;
- medicines used for HIV/AIDS (e.g. ritonavir, indinavir, nelfinavir);
- medicines used to treat fungal infections (e.g. ketoconazole, itraconazole);
- medicines used to treat bacterial infections (such as rifampicin, erythromycin, clarithromycin, telithromycin);
- verapamil, diltiazem (heart medicines);
- simvastatin (a medicine used to control high cholesterol levels);
- dantrolene (infusion for severe body temperature abnormalities);
- medicines used to protect against transplant rejection (ciclosporin).

Eslo Tan[®] Tablets with food and drink

Grapefruit juice and grapefruit should not be consumed by people who are taking Eslo Tan[®] Tablets. This is because grapefruit and grapefruit juice can lead to an increase in the blood levels of the active ingredient S(-)Amlodipine, which can cause an unpredictable increase in the blood pressure-lowering effect of S(-)Amlodipine.

Pregnancy, breast-feeding and fertility

Pregnancy

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Eslo Tan[®] before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Eslo Tan[®]. Eslo Tan[®] is not recommended in early pregnancy (first 3 months), and must not be taken when more than 3 months pregnant.

Breast-feeding

Tell your doctor if you are breast-feeding or about to start breast-feeding. Eslo Tan[®] is not recommended for mothers who are breastfeeding, and your doctor may choose another treatment for you if you wish to breastfeed, especially if your baby is a newborn or was born prematurely. Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

This medicine may make you feel dizzy. This can affect how well you can concentrate. So, if you are not sure how this medicine will affect you, do not drive, use machinery, or do other activities that you need to concentrate on.

3. How to take Eslo Tan[®] Tablets

Always take or give this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. This will help you get the best results and lower the risk of side effects.

The usual dose of Eslo Tan[®] is one tablet per day.

- It is preferable to take your medicine at the same time each day.
- Swallow the tablets with a glass of water.
- You can take Eslo Tan[®] with or without food. Do not take Eslo Tan[®] with grapefruit or grapefruit juice.

Patients with kidney or liver problems

You should tell your doctor if you have kidney or liver problems as your doctor may need to alter the normal dose or medicine.

If you take more Eslo Tan[®] than you should

If you have taken too many tablets of Eslo Tan[®], or if someone else has taken your tablets, consult a doctor immediately.

If you forget to take or give Eslo Tan[®]

If you forget to take this medicine, take it as soon as you remember. Then take your next dose at its usual time. However, if it is almost time for your next dose, skip the dose you missed. Do not take a double dose to make up for a forgotten tablet.

If you stop taking Eslo Tan[®]

Stopping your treatment with Eslo Tan[®] may cause your disease to get worse. Do not stop taking your medicine unless your doctor tells you to.

4. Possible side effects

This medicine can cause side effects like all medicines, although not everybody gets them.

Some side effects can be serious and need immediate medical attention: A few patients have experienced these serious side effects (may affect up to 1 in 1,000 people). If any of the following happens, tell your doctor straight away: Allergic reaction with symptoms such as rash, itching, swelling of face or lips or tongue, difficulty breathing, low blood pressure (feeling of faintness, light-headedness).

Other possible side effects of Eslo Tan[®]:

Common (may affect up to 1 in 10 people): blocked nose, sore throat and discomfort when swallowing; headache; swelling of arms, hands, legs, ankles or feet; tiredness; asthenia (weakness); redness and warm feeling of the face and/or neck.

Uncommon (may affect up to 1 in 100 people): Dizziness; nausea and abdominal pain; dry mouth; drowsiness, tingling or numbness of the hands or feet; vertigo; fast heart beat including palpitations; dizziness on standing up; cough; diarrhoea; constipation; skin rash, redness of the skin; joint swelling, back pain; pain in joints.

Rare (may affect up to 1 in 1,000 people): Feeling anxious; ringing in the ears (tinnitus); fainting; passing more urine than normal or feeling more of an urge to pass urine; inability to get or maintain an erection; sensation of heaviness; low blood pressure with symptoms such as dizziness, light-headedness; excessive sweating; skin rash all over your body; itching; muscle spasm.

If any of these affect you severely, tell your doctor.

Side effects reported with S(-)amlodipine or losartan alone and either not observed with Eslo Tan[®] or observed with a higher frequency than with Eslo Tan[®]:

S(-)Amlodipine

Consult a doctor immediately if you experience any of the following very rare, severe side effects

after taking this medicine:

- Sudden wheeziness, chest pain, shortness of breath or difficulty in breathing.
- Swelling of eyelids, face or lips.
- Swelling of the tongue and throat which causes great difficulty breathing.
- Severe skin reactions including intense skin rash, hives, reddening of the skin over your whole body, severe itching, blistering, peeling and swelling of the skin, inflammation of the mucous membranes (Stevens-Johnson Syndrome, toxic epidermal necrolysis) or other allergic reactions.
- Heart attack, abnormal heart beat.

- Inflamed pancreas, which may cause severe abdominal and back pain accompanied with feeling of being very unwell.

Losartan

Not known (frequency cannot be estimated from the available data): Decrease in red blood cells, fever, sore throat or mouth sores due to infections; spontaneous bleeding or bruising; high level of potassium in the blood; abnormal liver test results; decreased renal functions and severely decreased renal functions; swelling mainly of the face and the throat; muscle pain; rash, purplish-red spots; fever; itching; allergic reaction; blistering skin (sign of a condition called dermatitis bullous).

If you experience any of these, tell your doctor straight away.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly: Website: www.zuventus.co.in and click the “Safety Reporting” located on the top of the home page.

By reporting side effects, you can help provide more information on the safety of this medicine.

You can also report the side effect with the help of your treating physician.

5. How to store Eslo Tan®

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister.

Do not store above 30°C.

Store in the original package in order to protect from moisture.

Do not use any Eslo Tan® pack that is damaged or shows signs of tampering.

6. Contents of the pack and other information

What Eslo Tan® tablets contain

Eslo Tan® 2.5 mg/50 mg tablets

The active substances of Eslo Tan® are S(-)Amlodipine as s(-) Amlodipine besylate and Losartan potassium. Each tablet contains 2.5 mg S(-)Amlodipine and 50 mg Losartan.

What Eslo Tan® looks like and contents of the pack

Eslo Tan® is available in packs of 10 tablets in PVC/PVDC-Alu blister strips.

Marketing Authorisation Holder

Zuventus Healthcare Limited
Zuventus House, Plot Y2, CTS No.: 358/A2,
Near Nahur Railway Station,

Nahur (W), Mumbai, 400078 Maharashtra, India.

This leaflet was last revised in 06/2023.