

Package Leaflet: Information for the user

TroxipTM-OD

Troxipide Prolonged-release Tablets

Read all of this leaflet carefully before your child starts taking this medicine because it contains important information for you. –

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for your child. Do not pass it on to others. It may harm them, even if their signs of illness are the same as your child's.
- If your child gets any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

What is in this leaflet?

1. What TroxipTM-OD is and what it is used for?
2. What you need to know before you take TroxipTM-OD?
3. How to take TroxipTM-OD?
4. Possible side effects
5. How to store TroxipTM-OD?
6. Contents of the pack and other information

1. What TroxipTM-OD is and what it is used for?

TroxipTM-OD is a tablet composed of Troxipide prolonged release 300mg.

Troxipide is a mucou-protective, anti-ulcer and anti-inflammatory agent that helps protect your stomach and intestines. It promotes tissue repair by:

- Increasing blood flow to the stomach's protective layer
- Helping protect the stomach lining from damage.
- Increasing prostaglandins (which are beneficial molecules in the stomach)
- Regeneration of the stomach lining and helps restore its normal function.

TroxipTM-OD tablet is used for the treatment of patient with:

- Acute and Chronic Gastritis
- Stomach Ulcers

2. What you need to know before you take TroxipTM-OD Tablet?

Do not take TroxipTM-OD Tablet:

If you are allergic to Troxipide or any of the ingredients in this medication.

Warnings and Precautions:

- **Pregnancy and Breastfeeding:** Do not take Troxip™-OD if you are pregnant or breastfeeding unless your doctor decides it is necessary.
- **Children:** Safety in children has not been established.
- **Elderly:** Use with caution due to potential impairment of physiological functions.

Other medicines and Troxip™-OD Tablet:

No known drug interaction has been reported yet.

Driving and using machines

Troxip™-OD tablets have no influence on the ability to drive and use machines.

3. How to take Troxip™-OD Tablet?

Always take Troxip™-OD tablet exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Dosage in Adults:

The general recommended dosage is one tablet once daily. The usual treatment duration is 8-12 weeks.

How to Take:

- Swallow the tablet whole with water.
- Take it with or after food or with a glass of milk.
- Do not crush, break, or chew the tablet.

Children and adolescents:

Safety in children has not been established and therefore, Troxip™-OD should not be used in this age group

If you take more Troxip™-OD tablet than you should

No overdose has been reported; however, it is advisable to go to the nearest casualty department or contact your doctor immediately. Take the tablet carton with you.

If you forget to take Troxip™-OD tablet

If you miss a dose, take one as soon as you can. If it is almost time for your next dose, skip the missed dose, do not take two doses at the same time to make up for the missed dose. If you have missed several doses, tell your doctor.

If you stop taking Troxip™-OD tablets

Do not stop taking the tablets or reduce the dose without telling your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist

4. Possible side-effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Common side effects include:

- Constipation
- Diarrhea
- Abdominal swelling
- Stomach discomfort
- Abnormal liver function tests
- General malaise

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly:

Website: www.zuventus.co.in and click the tab “Safety Reporting” located on the top end of the home page.

Website link: <https://www.zuventus.co.in/drug-safety-reporting>

By reporting side effects, you can help provide more information on the safety of this medicine.

You can also report the side effect with the help of your treating physician.

5. How to store Troxip™-OD?

Keep this medicine out of the sight and reach of children
Store in a cool, dry & dark place. Protect from light.

6. Contents of the pack and other information

Pack size: A blister strip of 10 tablets

Each prolonged release tablet contains:

Troxipide 300 mg

Colour: Lake of sunset yellow & Titanium Dioxide IP

Marketing Authorisation Holder

Zuventus Healthcare Ltd.

Zuventus House, Plot Y2, CTS No: 358/A2,

Near Nahur Railway Station,

Nahur (West), Mumbai 400 078.

This leaflet was last revised in June 2024.

